arXiv:0801.1997v2 [math.RA] 27 Mar 2008

AN UPPER BOUND FOR THE LOWER CENTRAL SERIES QUOTIENTS OF A FREE ASSOCIATIVE ALGEBRA

GALYNA DOBROVOLSKA AND PAVEL ETINGOF

ABSTRACT. Feigin and Shoikhet conjectured in [FS] that successive quotients $B_m(A_n)$ of the lower central series filtration of a free associative algebra A_n have polynomial growth. In this paper we give a proof of this conjecture, using the structure of a representation of W_n, the Lie algebra of polynomial vector fields on $\mathbb{C}^n$, on $B_m(A_n)$ which was defined in [FS]. Moreover, we show that the number of squares in a Young diagram D corresponding to an irreducible W_n-module in the Jordan-Hölder series of $B_m(A_n)$ is bounded above by the integer $(m-1)^2+2[\frac{n-2}{2}](m-1)$, which allows us to confirm the structure of $B_3(A_3)$ conjectured in [FS].

1. INTRODUCTION

Let $A:=A_n$ be the free associative algebra with n generators over $\mathbb{C}$. Consider its lower central series as a Lie algebra, i.e., the Lie ideals $L_m(A)\subset A$ defined recursively by $L_1(A)=A, L_i(A)=[A,L_{i-1}(A)]$, and the corresponding associated graded Lie algebra $B(A):=\oplus_{i\geq 1}B_i(A)$, where $B_i(A)=L_i(A)/L_{i+1}(A)$.

Giving each of the generators of A degree 1, we define a grading on A and hence on each of the spaces $B_i(A)$. It is an interesting (and, in general, unsolved) problem to determine the Hilbert series of $B_i(A)$ for each i with respect to this grading. For $i=1$, this is easy since $B_1(A)$ is the space of cyclic words in n letters. It particular, one can easily see that $B_1(A)$ has exponential growth. So at first sight one might expect that the spaces $B_i(A)$ for $i>1$ also have exponential growth. However, computer experiments performed by Eric Rains in 2005 suggested that, to the contrary, the spaces $B_i(A)$ for $i>1$ should have polynomial growth.

This phenomenon was studied systematically by Feigin and Shoikhet in the paper [FS]. The first important observation of Feigin and Shoikhet is that the image Z of $A[[A,A],A]A$ in $B_1(A)=A/[A,A]$ is central in the Lie algebra $B(A)$, and therefore, it is natural to define the space $\bar{B}_1(A)=B_1(A)/Z$ and the graded Lie algebra $\bar{B}(A)=\bar{B}_1(A)\oplus\oplus_{i\geq 2}B_i(A)$. Their second important observation is that the graded Lie algebra $\bar{B}(A)$ carries a natural grading-preserving action of the Lie algebra W_n of polynomial vector fields in n variables (while there is no such action on $B_1(A)$), and the W_n-modules $\bar{B}_1(A)$ and $B_i(A)$, $i\geq 2$, admit a Jordan-Hölder series whose simple composition factors are irreducible W_n-modules $\mathcal{F}_D$ of tensor fields corresponding to Young diagrams D with at most n rows.

Feigin and Shoikhet in [FS] determined the exact structure of $\bar{B}_1(A)$ and $B_2(A)$ as representations of W_n, showing that they have polynomial growth, and conjectured that for any m, $B_m(A)$ has polynomial growth, i.e., the dimension $B_m(A_n)[\ell]$, the degree ℓ part of $B_m(A_n)$, grows like $c_{mn}\ell^{n-1}$, where c_{mn} is a constant.

In this paper we give a proof of this conjecture. In fact, we show that the W_n-modules $B_m(A_n)$ have finite length, which implies the conjecture. More specifically, we give an explicit upper bound on the number of squares in a Young diagram D if the corresponding tensor field module $\mathcal{F}_D$ over W_n occurs as a composition factor in $B_m(A_n)$. This bound not only implies that $B_m(A_n)$ has finite length, but also, when combined with the computation by Eric Rains (see [FS]), allows us to confirm the conjectural structure of $B_3(A_3)$ (as conjectured in [FS], it turns out to be isomorphic to $\mathcal{F}_{2,1,0}$).

Remark 1. In this paper we work over the ground field $\mathbb{C}$ following [FS], but the discussion carries over without changes to an arbitrary field of characteristic zero.

Acknowledgments. P.E. is grateful to B. Shoikhet for useful discussions. The work of P.E. was partially supported by the NSF grant DMS-0504847.

2. Results

2.1. **Tensor field modules over W_n.** The simple W_n-modules $\mathcal{F}_D$ are defined in the following way. Let D be a Young diagram with at most n rows. Then $\mathcal{F}_D$ is the irreducible W_n-submodule of the space of polynomial tensor fields on $\mathbb{C}^n$ of type D. In other words, we take a Young diagram D and the corresponding irreducible $\mathfrak{gl}(n,\mathbb{C})$-module F_D. Then we extend the action of $\mathfrak{gl}(n,\mathbb{C})$ on F_D to the action of W_n^0, the subalgebra of W_n of vector fields vanishing at the origin, so that quadratic and higher vector fields act by zero. Then, if D has more than one column, we let $\mathcal{F}_D = \mathrm{Hom}_{U(W_n^0)}(U(W_n), F_D)$ be the W_n-module coinduced from a $\mathfrak{gl}(n,\mathbb{C})$-module F_D, which is known to be irreducible. If D has one column, we take $\mathcal{F}_D$ to be the irreducible W_n-submodule of the coinduced module $\mathrm{Hom}_{U(W_n^0)}(U(W_n), F_D)$ (which is known to be unique in this case).

It is known that if D has more than one column, then $\mathcal{F}_D$ is the whole space of tensor fields, otherwise (if D has one column of k squares) $\mathcal{F}_D$ is the space of closed polynomial k-forms on $\mathbb{C}^n$. If $(k_1,\ldots,k_n)$ is a partition (possibly ending with some zeroes) corresponding to a Young diagram D, we will write $\mathcal{F}_{(k_1,\ldots,k_n)}$ to denote the module $\mathcal{F}_D$. For references about modules $\mathcal{F}_D$, see [FF] or [F]; for reference about Schur modules F_D, see [Ful].

2.2. **The main result.** The main result of this paper is the following theorem, proved in the next section.

Theorem 2. *For $m \geq 3,\ n \geq 2$ and $\mathcal{F}_D$ in the Jordan-Hölder series of $B_m(A_n)$, we have the following estimate on the size (i.e., the number of squares) of the Young diagram D:*

$$|D| \leq (m-1)^2 + 2[\frac{n-2}{2}](m-1)$$

(where $[x]$ denotes the integer part of x).

Corollary 3. *For all $m \geq 2, n \geq 2$, $B_m(A_n)$ has finite length as a W_n-module. In particular,* $\dim B_m(A_n)[\ell] \sim c_{mn}\ell^{n-1}$*, as $\ell \to \infty$, where $c_{mn} > 0$ is a constant.*

Proof. It suffices to consider the case $m \geq 3$, as the case $m = 2$ is considered in [FS]. The first statement follows from Theorem 2 and the fact that $\dim B_m(A_n)[\ell] < \infty$. The second statement follows from the first statement. □

Corollary 4. $B_3(A_3) = \mathcal{F}_{2,1,0}$.

Proof. According to the MAGMA computation by Eric Rains (see [FS], formula (17) for $H_3(u,t)$), the characters of both sides agree up to degree 6 inclusively (see [FS], section 4.2). So if in addition to $\mathcal{F}_{2,1,0}$, $B_3(A_3)$ had included another $\mathcal{F}_D$, then D would have to have at least 7 squares. But according to Theorem 2, D can have at most 4 squares. Thus, there is no additional constituents, and we are done. $\square$

Remark 5. In a similar way, one can rederive the results of [DKM] on the structure of $B_3(A_2)$ and $B_4(A_2)$. Namely, for $B_4(A_2)$, the bound for the size of D given by Theorem 2 is 9, while the Hilbert series of $B_4(A_2)$ is computed by Eric Rains using MAGMA up to degree 9 (see [FS]).

3. Proof of Theorem 2

3.1. **The map ψ and its kernel.** Consider the polynomial rings $\mathcal{O}_{mn} := \mathbb{C}[x_{ij}]_{1\le i\le n, 1\le j\le m}$. Below we write $A := A_n$. Let $\Omega^{even}(\mathbb{C}^n)$ be the space of even polynomial differential forms, and set $\Omega^{mn} := \Omega^{even}(\mathbb{C}^n)^{\otimes m}$. As an $\mathcal{O}_{mn}$-module, $\Omega^{mn} \cong \mathcal{O}_{mn} \otimes \Lambda^{mn}$, where $\Lambda^{mn} := \Lambda^{even}(\mathbb{C}^n)^{\otimes m}$.

Recall that by a result of [FS], we have a surjective map $\xi : \Omega^{even}(\mathbb{C}^n) \to \bar{B}_1(A)$, which descends to an isomorphism $\Omega^{even}(\mathbb{C}^n)/\Omega^{even}_{exact}(\mathbb{C}^n) \cong \bar{B}_1(A)$, which we will also denote by ξ. For any $m \ge 2$, this isomorphism gives rise to a map $\psi : \Omega^{mn} \to B_m(A)$ defined as a composition of two maps

$$\Omega^{mn} = \Omega^{even}(\mathbb{C}^n)^{\otimes m} \to \bar{B}_1(A)^{\otimes m} \to B_m(A),$$

$\omega_1 \otimes \cdots \otimes \omega_m \mapsto b_1 \otimes \cdots \otimes b_m \mapsto [[b_1, b_2], \ldots, b_m]$, where $b_i := \xi(\omega_i)$. The map ψ is surjective, since by the results of [FS], the direct sum $\bar{B}_1(A) \oplus \bigoplus_{m\ge 2} B_m(A)$ is a graded Lie algebra generated in degree 1.

Recall also from [FS] that we have a natural isomorphism $\eta : \Omega^{even>0}_{exact}(\mathbb{C}^n) \to B_2(A)$, and upon identification by ξ, η, the bracket map $\bar{B}_1(A)^{\otimes 2} \to B_2(A)$ reduces to the map $\omega_1 \otimes \omega_2 \to d\omega_1 \wedge d\omega_2$.

Therefore, we can factor the map ψ as

$$\Omega^{even}(\mathbb{C}^n)^{\otimes 2} \otimes \Omega^{even}(\mathbb{C}^n)^{\otimes(m-2)} \to B_2(A) \otimes \bar{B}_1(A)^{\otimes(m-2)} \to B_m(A)$$

using the map $\eta \circ \phi : \Omega^{even}(\mathbb{C}^n)^{\otimes 2} \to B_2(A)$, where $\phi : \omega_1 \otimes \omega_2 \mapsto d\omega_1 \wedge d\omega_2$.

Lemma 1. *For any $i, j \in [1, n]$ and forms $\omega_1, \omega_2 \in \Omega^{even}(\mathbb{C}^n)$, the element $(x_{i1} - x_{i2})(x_{j1} - x_{j2})\omega_1 \otimes \omega_2$ is in the kernel of the map ϕ. Consequently, the submodule $(x_{i1} - x_{i2})(x_{j1} - x_{j2})\Omega^{mn}$ is in the kernel of ψ.*

Proof. We have

$$\phi((x_{i1} - x_{i2})\omega_1 \otimes \omega_2) = \{d(x_{i1}\omega_1) \wedge d\omega_2 - d\omega_1 \wedge d(x_{i2}\omega_2)\}|_{x_{i1}=x_{i2}=x_i}$$
$$= dx_i \wedge \omega_1 \wedge d\omega_2 + x_i d\omega_1 \wedge d\omega_2 - d\omega_1 \wedge dx_i \wedge \omega_2 - d\omega_1 \wedge x_i d\omega_2$$

(since ω_1 is even)

$$= dx_i \wedge (\omega_1 \wedge d\omega_2 + d\omega_1 \wedge \omega_2).$$
$$= dx_i \wedge d(\omega_1 \wedge \omega_2).$$

Using this, we compute

$$\phi((x_{i1} - x_{i2})(x_{j1} - x_{j2})\omega_1 \otimes \omega_2) =$$
$$= \phi((x_{j1} - x_{j2})(x_{i1}\omega_1 \otimes \omega_2) - (x_{j1} - x_{j2})(\omega_1 \otimes x_{i2}\omega_2))$$
$$= dx_j \wedge d(x_i\omega_1 \wedge \omega_2) - dx_j \wedge d(\omega_1 \wedge x_i\omega_2) = 0.$$

Therefore, ϕ maps $(x_{i1}-x_{i2})(x_{j1}-x_{j2})\Omega^{even}(\mathbb{C}^n)^{\otimes 2}$ to zero. In particular, this implies that $(x_{i1}-x_{i2})(x_{j1}-x_{j2})\Omega^{mn}$ belongs to the kernel of the map ψ. □

Lemma 2. *Let L be a Lie algebra, and $b_i \in L$, $i=1,...,m$. For any $k=1,...,m$, the bracket $[[b_1,b_2],\dots,b_m]$ is a linear combination of the brackets of the form $[[b_k,b_{l_1}],\dots,b_{l_{m-1}}]$ where $(l_1,\dots,l_{m-1})$ is a permutation of $[1,m]\setminus\{k\}$.*

Proof. This lemma is well known, but we will give a proof, as it is very short. The proof is by induction in k. Indeed, for $k=1,2$ the statement is true. To go from $k-1$ to k, we notice that by the Jacobi identity

$$[[b_1,b_2],\dots,b_{k-1}],b_k] = [[b_1,b_2],\dots,b_k],b_{k-1}] + [[b_1,b_2],\dots],[b_{k-1},b_k]].$$

Putting $b'_{k-1} := b_k$ we have that the first bracket on the RHS can be expressed as a linear combination of brackets of the form $[[b_k,b_{l_1}],\dots,b_{l_{k-1}}]$ by the induction assumption. Similarly, in the second bracket on the RHS we put $b''_{k-1} := [b_{k-1},b_k]$ and use the induction assumption to express the second bracket as a combination of brackets of the form $[[b_k,b_{k-1}],\dots]$. Bracketing the LHS and RHS with $b_{k+1},\dots,b_m$, we obtain the desired result. □

Let I be the ideal in $\mathcal{O}_{mn}$ generated by the polynomials

$$\prod_{s=1}^{k-1}(x_{i_s s}-x_{i_s k})(x_{j_s s}-x_{j_s k}).$$

where $2\le k\le m$ and arbitrary $i_s, j_s \in [1,n]$ for $s\in[1,k]$.

Lemma 3. *The submodule $I\Omega^{mn}$ of Ω^{mn} is W_n-invariant, and is contained in the kernel of the map ψ.*

Proof. First we show that $I\Omega^{mn}$ is stable under the action of of W_n on Ω^{mn}. For this, it suffices to show that the ideal I is W_n-invariant. Let $f\frac{\partial}{\partial y_m}$ be some vector field in W_n. Then we have:

$$f\frac{\partial}{\partial y_m}x_{ij} = \begin{cases} 0 & \text{if } k\neq j \\ f_j & \text{if } m=i \end{cases}$$

where f_j denotes f where instead of the variables y_l we substitute x_{lj}. Therefore,

$$(f\frac{\partial}{\partial y_m})\prod_{s=1}^{k-1}(x_{i_s s}-x_{i_s k})(x_{j_s s}-x_{j_s k})$$

$$=\prod_{s=1}^{k-1}(x_{i_s s}-x_{i_s k})(x_{j_s s}-x_{j_s k})\sum_{s=1}^{k-1}(\frac{f_s-f_k}{x_{i_s s}-x_{i_s k}}+\frac{f_s-f_k}{x_{j_s s}-x_{j_s k}})$$

The last factor is clearly a polynomial, so the right hand side belongs to I. This implies that I is W_n-invariant, as desired.

Next we prove that the element $\prod_{s=1}^{k-1}(x_{i_s s}-x_{i_s k})(x_{j_s s}-x_{j_s k})\omega_1\otimes\cdots\otimes\omega_m$ goes to zero under the map ψ. Let $b_i=\xi(\omega_i)$. By Lemma 2, in $B_m(A)$ we have

$$[[b_1,b_2],\dots,b_m] = \sum_\sigma c_\sigma[[b_k,b_{\sigma(1)}],\dots,b_{\sigma(m-1)}],$$

where σ is a bijection from $[1,m-1]$ to $[1,m]\setminus\{k\}$. Therefore, we have that

$$\omega_1\otimes\cdots\otimes\omega_m - \sum_\sigma c_\sigma(\omega_k\otimes\omega_{\sigma(1)})\otimes\cdots\otimes\omega_{\sigma(m-1)}$$

is in the kernel of ψ.

But by Lemma 1 $(x_{i_{\sigma(1)}\sigma(1)} - x_{i_{\sigma(1)}k})(x_{j_{\sigma(1)}\sigma(1)} - x_{j_{\sigma(1)}k})(\omega_k \otimes \omega_{\sigma(1)}) \otimes \cdots \otimes \omega_{\sigma(m-1)}$ is in the kernel of ψ.

Therefore, $I\Omega^{mn}$ is in the kernel of ψ. $\square$

Set $r := 2[\frac{n-2}{2}](m-1)$. Note that the space $\Lambda(\mathbb{C}^n)$ is equipped with the natural grading by rank of exterior forms; hence so is the space Λ^{mn}. Consider the space $\Lambda^{mn}_{>r}$ spanned by homogeneous elements of degree $> r$ in Λ^{mn}.

Lemma 4. *The submodule $\mathcal{O}_{mn} \otimes \Lambda^{mn}_{>r}$ of Ω^{mn} is W_n-invariant, and is contained in the kernel of ψ.*

Proof. Since vector fields act as Lie derivative on the components of Ω^{mn}, they leave rank unchanged, so $\mathcal{O}_{mn} \otimes \Lambda_{>r}$ is stable under the W_n-action on Ω^{mn}.

Now we show that this submodule is annihilated by ψ. Elements of $\psi(\mathcal{O}_{mn} \otimes \Lambda^{mn}_{>r})$ are linear combinations of elements of the form $[[b_1, b_2] \dots b_m]$, $b_i = \xi(\omega_i)$, where $\sum \mathrm{rk}\omega_i > 2[\frac{n-2}{2}](m-1)$. By Lemma 2, for every $k \in [1, m]$ we have $[[b_1, b_2] \dots b_m] = \sum_\sigma c_\sigma [[b_k, b_{\sigma(1)}] \dots b_{\sigma(m-1)}]$.

Pick k such that ω_k has maximal rank among $\omega_1, \dots, \omega_m$. Then $\mathrm{rk}\omega_k + \mathrm{rk}\omega_{\sigma(1)} \geq \frac{1}{m-1} \sum \mathrm{rk}\omega_i > 2[\frac{n-2}{2}]$. But $\mathrm{rk}\omega_k + \mathrm{rk}\omega_{i_1}$ is even so $\mathrm{rk}\omega_k + \mathrm{rk}\omega_{i_1} \geq 2([\frac{n-2}{2}] + 1)$. So we have

$$\mathrm{rk}(d\omega_k \wedge d\omega_{\sigma(1)}) = \mathrm{rk}\omega_k + \mathrm{rk}\omega_{\sigma(1)} + 2 \geq 2([\frac{n-2}{2}] + 2) = 2[\frac{n+2}{2}] > n.$$

Thus $d\omega_k \wedge d\omega_{\sigma(1)} = 0$ in $\Omega^{even}(\mathbb{C}^n)$.

But $[b_k, b_{\sigma(1)}] = \eta(d\omega_k \wedge d\omega_{\sigma(1)})$ in $B_2(A)$. So $[b_k, b_{\sigma(1)}] = 0$, and thus we have $[[b_1, b_2] \dots b_m] = 0$. Therefore the submodule $\mathcal{O}_{mn} \otimes \Lambda^{mn}_{>r}$ is in the kernel of ψ. $\square$

3.2. **The structure of O_{mn}/I as a graded space.** Note that the algebra $\mathcal{O}_{mn}$ has a natural grading, which assigns degree 1 to each generator, and I is a graded ideal. Thus, $\mathcal{O}_{mn}/I$ is a graded algebra.

Lemma 5. *We have an isomorphism of graded vector spaces*

$$\mathcal{O}_{mn}/I \cong \mathbb{C}[y_1, \dots, y_n] \otimes \bigotimes_{k=2}^{m} \mathrm{Sym}^{2k-3}(\mathbb{C} + \sum_{i=1}^{n} \mathbb{C}y_i),$$

where y_i have degree 1.

Proof. Denote by $\mathrm{in}(I)$ the set of initial (highest) terms of polynomials in I with respect to the lexicographic monomial order $x_{ij} > x_{kl}$ iff $j > l$ or $j = l$ and $i > k$. By a theorem about Gröbner bases (Theorem 15.3 in [E]) monomials not in $\mathrm{in}(I)$ form a vector space basis of $\mathcal{O}_{mn}/I$. By the form of the generators of I we see that $\mathrm{in}(I)$ is generated by the monomials $\prod_{s=1}^{2k-2} x_{i_s k}$, where $k \geq 2$, $i_s \in [1, n]$ is arbitrary. So monomials not in $\mathrm{in}(I)$ are precisely the monomials in $\mathbb{C}[x_{11}, \dots, x_{n1}] \otimes \bigotimes_{k=2}^{m} \mathrm{Sym}^{2k-3}(\mathbb{C} + \sum_{i=1}^{n} \mathbb{C}x_{ik})$. $\square$

3.3. **Proof of Theorem 2.** For any graded space M define its character (the Hilbert series of M) $\mathrm{char}M = \sum_a \dim M[a] t^a$, where $M[a]$ is the a^{th} graded piece of M. For instance, $\mathrm{char}\mathcal{F}_D = \frac{P_D(t)}{(1-t)^n}$, where $P_D(t)$ is a polynomial of degree n if D has one column, and $P_D(t) = N_D t^{|D|}$ if D has more than one column, where N_D is the dimension of the irreducible representation of GL_n corresponding to D.

The space $\Omega^{mn} = \mathcal{O}_{mn} \otimes \Lambda^{mn}$ has a natural grading coming from the grading on the factors, and we have a surjective map of graded vector spaces $\psi : \Omega^{mn} \twoheadrightarrow B_m(A_n)$. By Lemma 3, $I\Omega^{mn} = I\mathcal{O}_{mn} \otimes \Lambda^{mn}$ is in the kernel of ψ, and by Lemma 4, $\mathcal{O}_{mn} \otimes \Lambda^{mn}_{>r}$ is in the kernel of ψ. So we have a surjective map

$$(\mathcal{O}_{mn} \otimes \Lambda^{mn})/(I\mathcal{O}_{mn} \otimes \Lambda^{mn} + \mathcal{O}_{mn} \otimes \Lambda^{mn}_{>r}) \twoheadrightarrow B_m(A_n).$$

Thus we have a surjective map

$$\hat{\psi} : (\mathcal{O}_{mn}/I) \otimes (\Lambda^{mn}/\Lambda^{mn}_{>r}) \twoheadrightarrow B_m(A_n),$$

which is a homomorphism of W_n-modules. We also notice that the space $\Lambda^{mn}/\Lambda^{mn}_{>r}$ is naturally identified with $\Lambda^{mn}_{\leq r}$.

Let $\mu(D)$ be the multiplicity of $\mathcal{F}_D$ in $(\mathcal{O}_{mn}/I) \otimes (\Lambda^{mn}/\Lambda^{mn}_{>r})$, and $\nu(D)$ be the multiplicity of $\mathcal{F}_D$ in $B_m(A_n)$. Since $\hat{\psi}$ is surjective, $\nu(D) \leq \mu(D)$. Also, we have

$$\sum_D \mu(D)\mathrm{char}\mathcal{F}_D = \mathrm{char}(\mathcal{O}_{mn}/I)\mathrm{char}(\Lambda^{mn}_{\leq r}). \tag{1}$$

From Lemma 5 we have

$$\begin{aligned}\mathrm{char}\mathcal{O}_{mn}/I &= \mathrm{char}\mathbb{C}[y_1, \dots, y_n] \prod_{k=2}^{m} \mathrm{char}\ \mathrm{Sym}^{2k-3}(\mathbb{C} + \sum_{i=1}^{n} \mathbb{C}y_i) \\ &= \frac{1}{(1-t)^n} \prod_{k=2}^{m} (\sum_{a=0}^{2k-3} \binom{a+n-1}{n} t^a) \\ &= \frac{1}{(1-t)^n}(c_0 t^{(m-1)^2} + \text{l.o.t.}),\end{aligned}$$

where $c_0 \neq 0$ is a constant and l.o.t. denote monomials of smaller degree than $(m-1)^2$. The character of $\Lambda^{mn}_{\leq r}$ is a polynomial of degree r, $C_0 t^r + \text{l.o.t.}$

Thus, multiplying (1) by $(1-t)^n$, we obtain

$$\sum_D \mu(D) P_D(t) = Q(t),$$

where $Q(t)$ is a polynomial of degree $(m-1)^2 + r$.

Now, for $m \geq 3, n \geq 2$

$$n \leq 2(n-1) \leq 4[n/2] = 4 + 4[(n-2)/2] \leq (m-1)^2 + r.$$

So, we get

$$\sum_D \mu(D) N_D t^{|D|} = Q_*(t),$$

where Q_* is another polynomial of degree $(m-1)^2 + r$. Hence $\mu(D) = 0$ for $|D| > (m-1)^2 + r$ and hence $\nu(D) = 0$ for $|D| > (m-1)^2 + r$, as desired.

REFERENCES

[DKM] G. Dobrovolska, J. Kim, X. Ma, On the lower central series of an associative algebra, arXiv:0709.1905.

[E] D.Eisenbud, Commutative algebra with a view toward algebraic geometry, Graduate texts in mathematics Vol.150, Springer (1995).

[FF] B. Feigin, D.B. Fuks, Cohomologies of Lie Groups and Lie Algebras, in A.L. Onishchik, E.B. Vinberg (Eds.), Lie Groups and Lie Algebras II, Springer 2000.

[FS] B. Feigin, B. Shoikhet, On $[A, A]/[A, [A, A]]$ and on a W_n-action on the consecutive commutators of free associative algebras, Math. Res. Lett. 14 (2007), no. 5, 781795.

[F] D.B. Fuks, Cohomology of Infinite-Dimensional Lie Algebras, Consultants Bureau 1986.

[Ful] W. Fulton, Young Tableaux, London Math. Soc. Student Texts 35, Cambridge Univ. Press 1997.

Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA 02139, U.S.A.
E-mail address: `galyna@mit.edu`

Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA 02139, U.S.A.
E-mail address: `etingof@math.mit.edu`